

**The Contribution of Instanton-Antiinstanton Fluctuations to
the Ground State Energy in Supersymmetric Quantum Mechanical Models**

Avinash Khare

Institute of Physics, Sachivalaya Marg,
Bhubaneswar 751005, India.

email: khare@iopb.res.in

IP-BBSR/84-20, October 1984, SLAC-PPF T8447

Abstract

I show that the method of finding if the ground state energy is nonzero or not by saturating the functional integral directly with instanton-anti-instanton type fluctuations is not reliable either for the double or for the triple well potential models in supersymmetric quantum mechanics.

In the last few years, supersymmetry (SUSY) has attracted lot of attention as it appears to cure the gauge hierarchy problem in the grand unified theories. If SUSY has anything to do with nature, it is clear that it must be a badly broken symmetry. There are reasons to believe that SUSY may be broken dynamically by nonperturbative fluctuations rather than spontaneously at the tree level. It may therefore be worthwhile to understand the mechanism responsible for dynamical SUSY breaking.

Sometime ago, Witten [1] conjectured that instantons may be responsible for the dynamical SUSY breaking. Since it is not very easy to study this question for field theoretic models in 3+1 dimensions, many workers have studied it in SUSY quantum mechanics. Witten and specially Salomonson and van Holten [2] have discussed the celebrated $x^2 - x^4$ double well potential which has one instanton and have shown that SUSY is indeed dynamically broken in this example. Later on, Cooper and Freedman [3] have shown that instantons are not necessary for the SUSY breaking by considering a x^4 -potential. However, the real surprise came when Abbott and Zakrzewski [4] showed that instantons are not even sufficient for SUSY breaking. In particular, they discussed the potential $x^2 - x^4 - x^6$ having symmetric triple well [5] and showed that even though there are two instantons in this model still SUSY is unbroken. This counter example made it clear that the connection between instantons and dynamical SUSY breaking, if any, is not that straight forward [6].

In a recent paper, Kaul and Mizrachi [7] have reexamined this question and offered an explanation as to why SUSY is dynamically broken for the $x^2 - x^4$ potential with double well while it is unbroken for the triple well case ($x^2 - x^4 - x^6$). These authors saturated the functional integral directly with instanton-antiinstanton type fluctuations and showed that whereas in the double well case the ground state energy is raised by these fluctuations, in the triple well case it is not. According to these authors, their formalism can be easily generalized to field theoretical models in higher dimensions. In view of these attractive features, it may be worthwhile to critically analyze this approach. In particular, one would like to know if the approach gives reliable result for any double or triple well potential model or not. Further, it would appear from the work of Kaul and Mizrachi [7] that whereas SUSY would be dynamically broken for any double well potential, it would remain unbroken for any triple well case. Is this conjecture really true ? Unfortunately, as I show below the answer to all these questions is “no”. In particular, I offer

concrete examples with double and triple well potentials and show that in both cases this method leads to wrong conclusion regarding the question of SUSY breaking. I also show that unbroken or broken SUSY has nothing to do with the triple or double well structure of the potential.

The Minkowskian action for a supersymmetric classical particle is given by

$$A_M = \frac{1}{2} \int dt [\dot{x}^2 - S^2(x) + i\psi^T \dot{\psi} - S'(x)\psi^T \sigma_2 \psi], \quad (1)$$

where $S'(x)$ denotes the derivative with respect to its argument x , $\dot{x}$ means derivative with respect to time t while ψ is a two-component anticommuting variable. Note that this action is invariant under the supersymmetric transformation

$$\delta_\varepsilon x = \varepsilon^T \sigma_2 \psi, \quad \delta_\varepsilon \psi = [-i\sigma_2 \dot{x} - S(x)]\varepsilon, \quad \delta_\varepsilon \psi^T = \varepsilon^T [i\sigma_2 \dot{x} - S(x)]. \quad (2)$$

Let us first consider the following double well potential

$$V \equiv \frac{1}{2} S^2 = \frac{a\lambda^2}{9m^2} (x^2 - \frac{\alpha^2 m^2}{\lambda})^2 (x^2 + \frac{\beta^2 m^2}{\lambda}), \quad (3)$$

where

$$\alpha^2 = \frac{3}{2a} [1 + (1 - \frac{2a}{3})^{1/2}], \quad \beta^2 = \frac{3}{a} [(1 - \frac{2a}{3})^{1/2} - \frac{1}{2}]. \quad (4)$$

while a is an arbitrary parameter. In order that V has a double well structure it is necessary that $0 < a < 9/8$. This ensures that β^2 is always positive. In fact, if one had chosen $a = 9/8$ then $\beta^2 = 0$ and the potential (3) reduces to the symmetric triple well of [7]. For $0 < a < 9/8$, the potential (3) has two classical ground states denoted by $-\pm >$ corresponding to $x = \pm \alpha m / \sqrt{\lambda}$ and $\psi = 0$. Note that for these ground states the classical vacuum energy is zero so that SUSY remains unbroken at the tree level. What about dynamical SUSY breaking because of nonperturbative effects? Since for this model, asymptotically $S \sim x^3$ hence a la Witten [1] it follows that SUSY remains unbroken in this model.

Let us now try to follow the arguments of [7] and see if by saturating the functional integral directly with instanton-antiinstanton type fluctuations one would arrive at the same conclusion or not. For the double well potential (3) there is one instanton and one antiinstanton solution given by [8] $\psi = 0$ and

$$x_I(t - t_1) = \frac{m}{\sqrt{\lambda}} (\frac{3\varepsilon}{a^2})^{1/2} (1 + \frac{9\varepsilon}{a^2})^{-1/4} \frac{\sinh y}{[1 + \frac{1}{3}(1 - [1 + \frac{9\varepsilon}{a^2}]^{-1/2}) \sinh^2 y]^{1/2}}, \quad (5)$$

$$x_{\overline{I}}(t - t_1) = -x_I(t - t_1). \quad (6)$$

Here

$$\varepsilon = \frac{1}{2}[(1 - a) + (1 - \frac{2a}{3})^{3/2}], \quad (7)$$

while

$$y = (1 + \frac{9\varepsilon}{a^2})^{1/4} m(t - t_1). \quad (8)$$

The classical action for both these solutions is the same [8]

$$A_0 = \frac{81m^3\varepsilon}{\sqrt{2}\lambda a^2} \frac{F(3, \frac{5}{2}, \frac{11}{4}; \gamma)\Gamma(11/4)}{(1 + \frac{9\varepsilon}{a^2})^{1/4}[1 - (1 + \frac{9\varepsilon}{a^2})^{-1/2}]^3\Gamma(5/2)\Gamma(1/4)} \quad (9)$$

where

$$\gamma = -\frac{[2(1 + \frac{9\varepsilon}{a^2})^{1/2} + 1]}{[(1 + \frac{9\varepsilon}{a^2})^{1/2} - 1]}. \quad (10)$$

Now one can essentially run through the arguments of [7]. As is well known, single instanton or antiinstanton will not contribute to the vacuum functional integral because of the zero modes of the relevant fermion determinant obtained by integrating over fermionic degrees of freedom. However, instanton-antiinstanton configurations will contribute since there are no exact fermionic zero modes for such configurations. By following through the arguments of [7] one can calculate the lowest order effect from such fluctuations and one finds that one has two ground states with equal energies given by

$$\langle + | H | + \rangle = \langle - | H | - \rangle \simeq \frac{m\hbar}{\sqrt{2}\pi} e^{-2A_0/\hbar}, \quad (11)$$

where A_0 is as given by eq. (9). Thus, according to this approach, to the lowest order, SUSY is dynamically broken which is contrary to the exact result that SUSY is unbroken for the potential (3). This counter example clearly demonstrates that the method of [7] is not reliable for double well potentials. Further it also shown that SUSY need not necessarily be broken for double well potentials.

It may be worthwhile to point out that for the potential (3) if one had instead followed the approach of [2] and calculated the single instanton induced vacuum expectation values of SUSY generators then also one would have arrived at the wrong conclusion that SUSY is dynamically broken for this model.

Let us now turn to the case of the triple well potentials and enquire about the reliability of the approach proposed in [7]. Let us consider the following triple well potential

$$V \equiv \frac{1}{2}S^2 = \frac{\lambda^3 x^2}{2m^4} \left(x^2 - \frac{m^2}{\lambda}\right)^2 \left(x^2 + \frac{m^2}{\lambda}\right). \quad (12)$$

The potential (12) has three classical ground states denoted by $| + \rangle$, $| - \rangle$ and $| 0 \rangle$ corresponding to $x = +m/\sqrt{\lambda}$, $-m/\sqrt{\lambda}$, 0 respectively (and $\psi = 0$). Note that the vacuum energy is zero for all these ground states so that SUSY is unbroken at the tree level. What about dynamical SUSY breaking? Since asymptotically $s \sim x^4$ hence a la Witten [1] it follows that the ground state energy $E_0 > 0$ and hence SUSY is dynamically broken in this model.

Let us now follow the approach of [7] and see if by saturating the functional integral directly with non-perturbative fluctuations one arrives at the same conclusion or not. For the triple well potential (12) it is clear that there will be two types of instantons which will interpolate between the classical ground states $x = 0$ and $x = \frac{m}{\sqrt{\lambda}}$ as well as $x = -\frac{m}{\sqrt{\lambda}}$ and $x = 0$ as t goes from $-\infty$ to $t \rightarrow \infty$ respectively. Similarly there will be two types of antiinstantons which will interpolate between the classical ground states $x = \frac{m}{\sqrt{\lambda}}$ and $x = 0$ as well as $x = 0$ and $x = -m/\sqrt{\lambda}$ as t goes from $-\infty$ to $t \rightarrow \infty$ respectively. The action for all these instantons and antiinstantons is the same:

$$A_0 = \frac{4m^3}{15\lambda}. \quad (13)$$

Single instanton or antiinstanton again does not contribute to the vacuum functional integral because of the fermionic zero modes. The possible next order contribution to the vacuum functional integral will again come from instanton-antiinstanton configurations. In this case, there will also be contributions from two instantons and two antiinstantons representing quantum tunneling from $| - \rangle$ to $| + \rangle$ and $| + \rangle$ to $| - \rangle$. One can calculate the lowest order effect from such vacuum fluctuations by following the approach of [7]. One would then find that to the lowest order

$$\langle 0 | H | 0 \rangle_{x_I^{(+)} x_T^{(+)}} = \langle 0 | H | 0 \rangle_{x_I^{(-)} x_T^{(-)}} \simeq \frac{m\hbar}{\sqrt{2\pi}} e^{-2A_0/\hbar} = X, \quad (14)$$

so that

$$\langle 0 | H | 0 \rangle_{x_I^{(+)} x_T^{(+)} + x_I^{(-)} x_T^{(-)}} = 2X. \quad (15)$$

Further

$$\begin{aligned}
& \langle + | H | + \rangle_{x_I^{(+)} x_I^{(-)}} = \langle - | H | - \rangle_{x_I^{(-)} x_I^{(-)}} = \langle + | H | - \rangle_{x_I^{(+)} x_I^{(-)}} \\
& = \langle - | H | + \rangle_{x_I^{(+)} x_I^{(-)}} \simeq \frac{m\hbar}{\sqrt{2\pi}} e^{-2A_0/\hbar} = X,
\end{aligned} \tag{16}$$

to the same order. Here A_0 is as given by eq.(13). Hence to the lowest order, the Hamiltonian matrix for the low-lying states can be written as

$$H = \begin{pmatrix} x & 0 & x \\ 0 & 2x & 0 \\ x & 0 & x \end{pmatrix}. \tag{17}$$

The matrix has a zero eigenvalue and two equal non-zero eigenvalues. Thus according to the approach of [7], to the lowest order, SUSY is not dynamically broken in this model which is contrary to the exact result. Thus the approach of [7] does not give reliable answers for the triple well potentials either. Further, this counter example also demonstrates that SUSY need not necessarily be unbroken for triple well potentials.

Summarizing, it appears that the approach of [7] at least in its present form, does not give reliable answers for either the double or the triple well potentials. Perhaps, it is not enough to calculate the contributions from widely separated instantons and antiinstanton but one must also include the contribution from their mutual interactions. Clearly much more work is necessary in order to clarify the connection between instantons and dynamical SUSY breaking.

References

- [1] E. Witten, Nucl. Phys. **B 188** (1981) 513.
- [2] P. Salomonson and J.W. van Holten, Nucl. Phys. **B196** (1982) 509.
- [3] F. Cooper and B. Freedman, Ann. of Phys. (N.Y) **146** (1984) 262.
- [4] R.B. Abbott and W.J. Zakrzewski, Z. Physik **C20** (1984) 227.
- [5] A. Khare, Lett. Math. Phys. **3** (1979) 475.
- [6] A. Khare and J. Maharana, Z. Physik **C23** (1984) 791.
- [7] R.K. Kaul and L. Mizrachi, CERN preprint CERN-TH. 3944/84, July (1984).
- [8] S.N. Behera and A. Khare, Pramana **15** (1980) 245;
A. Khare and S.N. Behera, Pramana **14** (1980) 327.